

Osteoarthritis and Cartilage



Brief Report

Varus malalignment negates the structure-modifying benefits of doxycycline in obese women with knee osteoarthritis

S.A. Mazzuca^{†,*}, K.D. Brandt[‡], R. Chakr[†], K.A. Lane[†]

[†] Indiana University School of Medicine, Indianapolis, IN, USA

[‡] Kansas University Medical Center, Kansas City, KS, USA

ARTICLE INFO

Article history:

Received 7 October 2009

Accepted 19 March 2010

Keywords:

Structure-modification

Knee osteoarthritis

Doxycycline

Varus malalignment

SUMMARY

Objective: To estimate the extent to which varus malalignment, a source of abnormal intra-articular stresses in the medial tibiofemoral compartment and risk factor for progression of knee osteoarthritis (OA), may have diminished the structure-modifying benefit of doxycycline in knee OA.

Methods: Post hoc treatment group comparisons from a randomized, placebo-controlled trial of the effect of doxycycline (100 mg, twice daily) on medial joint space narrowing (JSN) in subgroups of varus and non-varus OA knees. Subjects ($N = 379$ with X-ray follow-up) were obese 45–64-year-old women with unilateral knee OA at baseline. JSN was measured manually in semiflexed anteroposterior (AP) radiographs acquired with standardized fluoroscopic positioning. The anatomic-axis angle (AAA) was measured in each baseline radiograph and transformed to an estimate of the mechanical-axis angle (MAA_{est}) using a validated regression equation. Knees with MAA_{est} $< 178^\circ$ were classified as varus.

Results: In our original comparison with placebo, doxycycline slowed the rate of medial JSN in OA knees by 38% at 16 months and by 33% at 30 months. Among non-varus OA knees, 16-month JSN in the doxycycline group was 44% slower than in the placebo group (0.09 vs 0.16 mm/year, $P = 0.080$), and 39% slower at month 30 (0.10 vs 0.17 mm/year, $P = 0.026$). JSN in varus knees (0.20–0.27 mm/year) was more rapid than in non-varus knees ($P = 0.083$) and unaffected by doxycycline.

Conclusion: Varus malalignment negated the slowing of structural progression of medial-compartment OA by doxycycline. To our knowledge, this is the first report documenting that static varus angulation can negate a pharmacologic structure-modifying effect.

© 2010 Osteoarthritis Research Society International. Published by Elsevier Ltd. All rights reserved.

Introduction

While recent reviews have highlighted methodological advances that have increased the feasibility of clinical trials of disease-modifying OA drugs (DMOADs)^{1–3}, the results of randomized controlled trials (RCTs) of drugs intended to slow the structural damage of OA have been inconsistent. One reason for this inconsistency may be that pharmacotherapy intended to slow the deterioration of articular cartilage does nothing to negate the effects of abnormal intra-articular stresses that are the prime force in the etiopathogenesis and progression of common garden-variety OA⁴.

We have shown that, compared to placebo, treatment with doxycycline slowed the rate of radiographic joint space narrowing (JSN) in the medial-compartment of knees with established OA⁵.

Because varus malalignment increases loading of the medial tibiofemoral compartment⁶ and, hence, the risk of OA progression⁷, it may have attenuated the pharmacologic benefit of doxycycline. Herein we describe results of subgroup analyses from this trial to ascertain whether varus malalignment decreased the structure-modifying effect of doxycycline in knee OA.

Methods

All procedures for this RCT, including written informed consent, were approved by the Institutional Review Board (IRB) of Indiana University Purdue University Indianapolis, where the coordinating Center was located and by IRBs at other participating institutions⁵.

Subjects

The original sample of the doxycycline RCT consisted of 431 women recruited at six clinical centers in the United States. All were 45–64 years of age, in the upper tertile of sex-, age- and race-based norms for body mass index (BMI) and had unilateral knee OA

* Address correspondence and reprint requests to: Steven A. Mazzuca, Indiana University School of Medicine, Department of Medicine, Rheumatology Division, Clinical Building Room 370, 541 Clinical Drive, Indianapolis, IN 46202, USA. Tel: 1-317-274-0676; Fax: 1-317-274-7792

E-mail address: smazzuca@iupui.edu (S.A. Mazzuca).

(grade 2 or 3 by Kellgren and Lawrence (K&L) criteria⁸) at baseline. Data for the present analysis were derived from 379 subjects (88% of those randomized) who underwent the interim (16-month) radiographic examination and 361 subjects (84% of those randomized) who completed the closeout (30-month) assessment.

Measurement of JSN

The primary outcome of the RCT was radiographic JSN in the medial tibiofemoral compartment. As previously reported⁵, semi-flexed anteroposterior (AP) knee radiographs were acquired at baseline and at months 16 and 30 according to Buckland–Wright's protocol for fluoroscopically standardized radioanatomic positioning of the knee⁹. Quantitative JSN estimates were based on serial magnification-corrected measurements of minimum joint space width (JSW) obtained with a screw-adjustable calipers and graduated loupe according to the method described by Lequesne¹⁰.

Measurement of malalignment

The anatomic-axis angle (AAA) in the frontal plane was measured by one of two readers (RC, SAM) in each baseline radiograph of the index (OA) knee, following the methods described by Kraus *et al.*¹¹. AAA was measured by goniometer along the medial aspect of the lower extremity. The AAA was formed by lines that began at points bisecting the widths of the proximal femur and distal tibia (~10 cm from the joint space) and converged at a point midway between the tibial spines. Inter-reader reproducibility (intra-class correlation) of AAA measurements from 36 randomly selected radiographs was 0.95.

Measurements of AAA were transformed into estimates of the mechanical-axis angle (MAA_{est}) by use of the regression equation ($MAA_{est} = 0.915 \times AAA + 13.895$, $R^2 = 0.77$) validated by Hinman *et al.*¹². In their study, MAA estimated from AAA measured in knee radiographs accounted for 77% of variation in actual MAA measured in concurrent long-limb radiographs. Knees in the present study with $MAA_{est} < 178^\circ$ were classified as exhibiting varus angulation, by norms established by Moreland *et al.*¹³.

Analysis

Separate treatment group comparisons were performed on subgroups of varus and non-varus OA knees using the same mixed-effect (repeated measures) linear model employed in the original analysis of doxycycline effects⁵. To test whether the rates of medial-compartment JSN in varus and non-varus subgroups were similar, we employed an additional repeated measures model that included all knees and a dichotomous indicator of varus angulation ($MAA_{est} < 178^\circ$ vs $\geq 178^\circ$) as a covariate. Terms were included in each model to account for interactions between treatment group and visit, clinical center and varus malalignment (third model only). All comparisons were adjusted for baseline knee pain and JSW, visit (interim or closeout) and clinical center.

Results

Of the 379 subjects who underwent the interim (month 16) radiographic examination, the index (OA) knee exhibited varus angulation at baseline in 82 (22%). At closeout, 77 of the 361 index knees that were available for analysis (21%) had exhibited varus angulation in the baseline radiograph. As described previously⁵, all subjects enrolled in the RCT were obese, middle-aged women with unilateral K&L grades 2–3 OA at baseline. Subgroups of subjects with varus and non-varus index knees were similar with respect to age, BMI, race and baseline knee pain. However, varus knees were significantly more likely than

non-varus knees to have K&L grade 3 OA at baseline (64% vs 32%, $P < 0.0001$). Consistent with definitions of OA severity by K&L criteria⁸, mean JSW in varus knees was significantly smaller at baseline than that in non-varus knees (2.77 vs 3.90 mm, $P < 0.0001$).

Table 1 contains a summary of non-annualized JSN in the doxycycline and placebo treatment arms among subgroups of varus and non-varus OA knees. In both treatment groups, over both intervals, varus knees exhibited a greater loss of JSW than non-varus knees ($P = 0.083$ after adjustment for treatment group, baseline knee pain and JSW, visit, clinical center, and interactions between treatment group and both visit and clinical center).

Notably, group comparisons among varus knees showed similar rates of JSN in the two treatment groups, as reflected in average loss of minimum medial JSW at both month 16 (0.26 vs 0.36 mm, $P = 0.230$) and month 30 (0.49 vs 0.55 mm, $P = 0.448$). In contrast, among non-varus knees, the difference between treatment groups with respect to mean JSN favored doxycycline at both intervals. The mean difference between treatment groups was marginally significant as early as month 16 (0.12 vs 0.21 mm, $P = 0.080$) and unequivocally significant at month 30 (0.26 vs 0.42 mm, $P = 0.026$).

Discussion

Knee malalignment was not an exclusion criterion for the doxycycline trial, which we initiated in 1996, or for many other contemporaneous DMOAD trials in knee OA. Of eight knee RCTs of DMOADs that we reviewed previously³, only one¹⁴ excluded subjects with “clinically important” axial deviation of the lower extremity. Although current protocols for RCTs of putative DMOADs now often exclude subjects with significant varus or valgus deformity on the presumption that the efficacy of the drug may be diminished in such patients, to our knowledge, this is the first report documenting that static varus angulation can negate a pharmacologic structure-modifying effect.

In our original comparison of outcomes in active treatment and placebo groups⁵, oral doxycycline, 100 mg bid, slowed the annualized rate of medial JSN in OA knees by 38% at 16 months ($P = 0.027$) and by 33% at 30 months ($P = 0.017$). The *post hoc* subgroup analyses in the present study indicate, however, that doxycycline had no significant effect on the comparatively rapid rate of JSN in the medial tibiofemoral compartment of the varus knee (0.20–0.27 mm/year). These results are in marked contrast to the outcomes in non-varus knees, in which doxycycline slowed JSN by 43% at month 16, and by 38% at month 30, relative to placebo (Table 1).

Our decision to combine for analysis knees with neutral or valgus angulation at baseline into a “non-varus” subgroup was intended to maximize the power of treatment group comparisons to elucidate how varus malalignment influenced the effect of doxycycline on the primary outcome, JSN in the medial tibiofemoral compartment.

Table 1

Non-annualized medial-compartment JSN (mean \pm SD, mm) by treatment group: results in varus and non-varus OA knees

	Varus OA knees ($MAA_{est} < 178^\circ$)			Non-varus OA knees ($MAA_{est} \geq 178^\circ$)		
	Doxycycline	Placebo	P^*	Doxycycline	Placebo	P^*
16-Month JSN	0.26 \pm 0.39	0.36 \pm 0.57	0.230	0.12 \pm 0.42	0.21 \pm 0.52	0.080
N of knees	37	45		151	146	
30-Month JSN	0.49 \pm 0.64	0.55 \pm 0.62	0.448	0.26 \pm 0.58	0.42 \pm 0.73	0.026
N of knees	35	42		146	138	

Note: estimated MAA_{est} was based on measurement of the AAA¹².

* From mixed-effect (repeated measures) linear models adjusted for treatment group, baseline knee pain and JSW, visit (interim or closeout), clinical center, treatment group–clinical center interaction and treatment group–visit interaction.

However, because of its propensity toward lateral progression of OA and a corresponding pseudo-widening of medial joint space, the valgus knee may have been ill-suited for the demonstration of a DMOAD effect on progression of JSN in the medial-compartment. To explore the extent to which inclusion of subjects with valgus knees in the doxycycline RCT made it more difficult to demonstrate a significant DMOAD effect, we examined the annualized rate of medial-compartment JSN over 30 months (mean \pm SE) in OA knees with varus, neutral or valgus angulation at baseline (Fig. 1). Consistent with the results shown in Table I, at 30 months the rate of JSN in the 35 varus knees of subjects treated with doxycycline was only 11% slower than that in 42 varus knees of subjects who received placebo. In contrast, in neutrally aligned knees ($N = 37$ and 49 , respectively), doxycycline slowed the annual rate of medial JSN by nearly 50%, compared to placebo. Two findings are apparent in data from knees with valgus angulation ($MAA_{est} \geq 181^\circ$) at baseline: the first is that, not surprisingly, the rate of medial JSN was much slower in valgus knees than in either varus or neutral knees. Second, despite less rapid JSN in the medial-compartment, the mean rate of JSN in the 109 valgus knees of subjects treated with doxycycline was 25% slower than that in the 89 valgus knees of those receiving placebo.

These data suggest that while the rapid medial-compartment JSN in varus knees was unaltered by treatment with doxycycline, the drug retarded JSN in valgus knees – although to an extent only about half as great as in neutrally aligned knees. Inclusion of valgus knees in the RCT did not preclude detection of a significant protective effect of doxycycline against medial JSN in either our original analysis of all OA knees⁵ or in the current analysis of the non-varus subgroup.

This *post hoc* analysis has some limitations: first, restriction of enrollment in the doxycycline RCT to obese women may have limited the generalizability of any conclusions about the benefits of doxycycline as a DMOAD in the OA population at large. However, this narrow focus of the RCT did not compromise its value as a demonstration of the concept of structure-modification in knee OA – or of the present subgroup analysis as an exploration of the possible moderating effects of malalignment.

Second, the percentage of knees that were classified as having varus angulation at baseline (21–22%) was smaller than expected and may have limited our power to detect treatment effects in this

subgroup. This low percentage is likely due, in part, to our exclusion of subjects with K&L grade 4 knee OA at baseline. In addition, a skew in the distribution of MAA_{est} , away from the varus and toward the valgus end of the spectrum, may be a limitation inherent in the measurement of AAA with short (10–12 cm) segments of the tibial and femoral shafts¹⁵ or a bias in the arithmetic transformation of AAA measurements into estimates of MAA ¹². However, the small (9%) difference in 30-month JSN between varus knees assigned to the doxycycline or placebo groups (Fig. 1) suggests that the statistical power of treatment group comparisons among knees classified as varus in this analysis was not a significant limitation.

Third, malalignment itself is only a crude indicator of an adverse biomechanical environment, in which abnormal intra-articular stress drives the etiopathogenetic mechanisms that underlie joint breakdown in OA and may counteract attempts at physiologic or pharmacologic repair⁴. The evidence argues that it is not varus *per se*, or obesity, *per se*, that is detrimental to the knee, but the consequences of the resulting alteration in mechanical loading for the articular cartilage and subchondral bone. However, a recent report suggesting that leptin, which is produced by adipose tissue, protected against development of knee OA in markedly obese mice¹⁶, suggests the story may be more complicated.

Varus–valgus malalignment, as measured in a full-limb radiograph or estimated in a knee film, is an indication of *static* malalignment, whereas a varus thrust visualized during gait reflects *dynamic* malalignment. Varus–valgus alignment is a key determinant of the peak adductor moment, a measure of the magnitude of the intrinsic compressive load on the medial tibiofemoral compartment in stance. Varus further increases the medial-compartment load during gait; valgus increases stress in the lateral compartment¹⁷. Miyazaki *et al.*¹⁸ found that the peak adductor moment predicted X-ray progression in subjects with medial-compartment OA.

Chang *et al.*¹⁹ reported that a visualized varus thrust was associated with a four-fold increase in the likelihood of progression of medial tibiofemoral compartment OA. Among varus knees, the presence of a visualized thrust further increased the odds of progression of medial-compartment JSN three-fold. Thus, among OA knees with static varus malalignment, a subset that is characterized by the presence of a thrust appears to be at particularly high risk for OA progression. However, a thrust effect on progression (i.e., JSN) was apparent in knees with relatively mild varus alignment ($<5^\circ$) but not in those with more severe varus. Although approximately 70% of the total force across the normal knee has been shown to act on the medial-compartment during level walking²⁰, Chang *et al.*¹⁹ considered that with increasing varus malalignment, nearly 100% of the load is transmitted medially and hence, in the presence of higher degrees of varus, thrust has little additional impact. In any event, the possibility should be considered that dynamic, as well as static, loading of the knee might mitigate the structure-modifying effect of a DMOAD.

The significance of the present findings for the design of clinical trials of DMOADs should be viewed in light of published evidence²¹ that argues powerfully that the underlying problem in common, garden-variety OA is a quantitative or qualitative abnormality in intra-articular stress, due to a variety of causes; and that if the abnormal stress is eliminated, the OA joint can heal, with marked structural and symptomatic improvement. We have suggested that if the abnormal stress on the OA joint is *not* corrected, the likelihood is small that a putative DMOAD will be efficacious²¹. Our present data support that earlier suggestion. We recommend that future clinical trials of pharmacologic or non-pharmacologic therapies aimed at disease modification control the extraneous effects of biomechanical factors (either through exclusion criteria or through measurement and stratified allocation to treatment




Fig. 1. Mean rate (\pm SE) of medial-compartment JSN over 30 months in varus, neutral and valgus knees: comparisons of knees treated with doxycycline and placebo. For varus, neutral and valgus knees, N s in the doxycycline treatment group were 35, 37 and 109, respectively. For the placebo arm of the RCT, the corresponding N s were 42, 49 and 89, respectively.

groups). Furthermore, we suggest that DMOAD effects may be attenuated also by genetic or developmental abnormalities in joint shape^{22,23} and by neuromuscular abnormalities (e.g., sarcopenia, proprioceptive defects) that impair micro-coordination and, like malalignment, can increase intra-articular stress.

Conflict of interest

The authors have no conflict of interest.

Acknowledgment

This work was supported in part by grants from NIH/NIAMS (R01 AR044370 and R01 AR052740).

References

- Hunter DJ, Hellio Le Graverand-Gastineau MP. How close are we to having structure-modifying drugs available? *Med Clin North Am* 2009;93:223–34.
- Qvist P, Bay-Jensen AC, Christiansen C, Dam EB, Pastoureau P, Karsdal MA. The disease modifying osteoarthritis drug (DMOAD): is it in the horizon? *Pharmacol Res* 2008;58:1–7.
- Brandt KD, Mazzuca SA. Lessons learned from nine clinical trials of disease-modifying osteoarthritis drugs. *Arthritis Rheum* 2005;52:3349–59.
- Brandt KD, Dieppe P, Radin E. Commentary: is it useful to subset “primary” OA? A critique based on evidence regarding the etiopathogenesis of osteoarthritis. *Semin Arthritis Rheum* 2009;39:81–95.
- Brandt KD, Mazzuca SA, Katz BP, Lane KA, Buckwalter KA, Yocum DE, et al. Effect of doxycycline on progression of osteoarthritis. *Arthritis Rheum* 2005;52:2015–25.
- Tetsworth K, Paley D. Malalignment and degenerative arthropathy. *Orthop Clin North Am* 1994;25:367–77.
- Sharma L, Song J, Felson DT, Cahue S, Shamiyeh E, Dunlop D. The role of knee alignment in disease progression and functional decline in knee osteoarthritis. *JAMA* 2001;286:188–95.
- Kellgren JH, Lawrence JS. Radiographic assessment of osteoarthritis. *Ann Rheum Dis* 1957;16:494–502.
- Buckland-Wright JC, Macfarlane DG, Williams SA, Ward RJ. Accuracy and precision of joint space width measurements in standard and macroradiographs of osteoarthritic knees. *Ann Rheum Dis* 1995;54:872–80.
- Lequesne M. Quantitative measurements of joint space during progression of osteoarthritis: chondrometry. In: Kuettner K, Goldberg V, Eds. *Osteoarthritic disorders*. Rosemont, IL: American Academy of Orthopedic Surgeons; 1995:427–44.
- Kraus VB, Vail TP, Worrell T, McDaniel G. A comparative assessment of alignment angle of the knee by radiographic and physical examination methods. *Arthritis Rheum* 2005;52:1730–5.
- Hinman RS, May RL, Crossley KM. Is there an alternative to the full-leg radiograph for determining knee joint alignment in osteoarthritis? *Arthritis Rheum* 2006;55:306–13.
- Moreland J, Bassett L, Hanks G. Radiographic analysis of the axial alignment of the lower extremity. *J Bone Joint Surg Am* 1987;69:745–9.
- Jubb RW, Piva S, Beinat L, Dacre J, Gishen P. A one-year, randomised, placebo (saline) controlled clinical trial of 500–730 kDa sodium hyaluronate (Hyalgan®) on the radiological change in osteoarthritis of the knee. *Int J Clin Pract* 2003;57:467–74.
- Sheehy L, Felson D, Cooke T. Is the femoral shaft – tibial shaft angle a reliable substitute for the hip–knee–ankle angle in knee osteoarthritis? *Osteoarthritis Cartilage* 2009;17:393.
- Griffin TM, Hueber JL, Kraus VB, Guilak F. Extreme obesity due to impaired leptin signaling in mice does not cause knee osteoarthritis. *Arthritis Rheum* 2009;60:2935–44.
- Sharma L, Dunlop DD, Cahue S, Song J, Hayes KW. Quadriceps strength and osteoarthritis progression in malaligned and lax knees. *Ann Intern Med* 2003;138:613–9.
- Miyazaki T, Wada M, Kawahara H, Sato M, Baba H, Shimada S. Dynamic load at baseline can predict radiographic disease progression in medial compartment knee osteoarthritis. *Ann Rheum Dis* 2002;61:617–22.
- Chang A, Hayes K, Dunlop D, Hurwitz D, Song J, Cahue S, et al. Thrust during ambulation and the progression of knee osteoarthritis. *Arthritis Rheum* 2004;50:3897–903.
- Schipplein OD, Andriacchi TP. Interaction between active and passive knee stabilizers during level walking. *J Orthop Res* 1991;9:113–9.
- Brandt KD, Dieppe PA, Radin EL. Etiopathogenesis of osteoarthritis. *Med Clin N Amer*. 2009;93:1–24.
- Doherty M, Courtney P, Doherty S, Jenkins W, Maciewicz R, Muir K, et al. Nonspherical femoral head shape (pistol grip deformity), neck shaft angle, and risk of hip osteoarthritis. *Arthritis Rheum* 2008;58:3172–82.
- Cooke D, Scudamore A, Li J, Wyss U, Bryant T, Costigan P. Axial lower-limb alignment: comparison of knee geometry in normal volunteers and osteoarthritis patients. *Osteoarthritis Cartilage* 1997;5:39–47.